



Geometric-Arithmetic Reverse and Sum Connectivity Reverse Indices of Silicate and Hexagonal Networks

Research Article

V.R.Kulli^{1,*}

1 Department of Mathematics, Gulbarga University, Gulbarga, Karnataka, India.

Abstract: We introduce a new index known as geometric-arithmetic reverse index of a molecular graph. In this paper, we compute geometric-arithmetic reverse index and sum connectivity reverse index of different chemically interesting networks like silicate networks and hexagonal networks.

MSC: 05C05, 05C12, 05C35.

Keywords: Geometric-arithmetic reverse index, sum connectivity reverse index, silicate network, hexagonal network.

© JS Publication.

1. Introduction

Let G be a finite, simple, connected graph with vertex set $V(G)$ and edge set $E(G)$. Let $d_G(v)$ denote the degree of a vertex v in G . Let $\Delta(G)$ denote the maximum degree among the vertices of G . The reverse vertex degree of a vertex u in G is defined as $c_u = \Delta(G) - d_G(u) + 1$. The reverse edge connecting the reverse vertices u and v will be denoted by uv . For all further notation and terminology we refer to reader to [1]. Chemical graph theory is a branch of Mathematical Chemistry which has an important effect on the development of the chemical sciences. A topological index is a numerical parameter mathematically derived from the graph structure. Numerous such topological indices have been considered in Theoretical Chemistry. Recently we introduced the atom bond connectivity reverse index [2] of a graph G as

$$ABCC(G) = \sum_{uv \in E(G)} \sqrt{\frac{c_u + c_v - 2}{c_u c_v}}.$$

The sum connectivity reverse index was introduced by Kulli in [3]. The sum connectivity reverse index of a graph G is defined as

$$SC(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{c_u + c_v}}.$$

Recently some reverse indices were studied, for example, in [4–6]. We now introduce the geometric-arithmetic reverse index of a graph G as

$$GAC(G) = \sum_{uv \in E(G)} \frac{2\sqrt{c_u c_v}}{c_u + c_v}. \quad (1)$$

Recently several topological indices were studied, for example, in [7–17]. In this paper, the geometric-arithmetic reverse index and sum connectivity reverse index of silicate networks and hexagonal networks are computed. For silicate networks and hexagonal networks see [18].

* E-mail: vrkulli@gmail.com

2. Results for Silicate Networks

Silicates are obtained by fusing metal oxide or metal carbonates with sand. A silicate network is symbolized by SL_n where n is the number of hexagons between the center and boundary of SL_n . A 2-dimensional silicate network is depicted in Figure 1.




Figure 1: A 2-dimensional silicate network

Let G be the graph of silicate network SL_n . From Figure 1, it is easy to see that the vertices of SL_n are either of degree 3 or 6. Then $\Delta(G) = 6$. By algebraic method, we obtain that $|V(SL_n)| = 15n^2 + 3n$ and $|E(SL_n)| = 36n^2$. In SL_n , by algebraic method, there are three types of edges based on the degree of the end vertices of each edge as follows:

$$\begin{aligned} E_{33} &= \{uv \in E(G) | d_G(u) = d_G(v) = 3\}, & |E_{33}| &= 6n. \\ E_{36} &= \{uv \in E(G) | d_G(u) = 3, d_G(v) = 6\}, & |E_{36}| &= 18n^2 + 6n. \\ E_{66} &= \{uv \in E(G) | d_G(u) = d_G(v) = 6\}, & |E_{66}| &= 18n^2 - 12n. \end{aligned}$$

We have $c_u = \Delta(G) - d_G(u) + 1 = 7 - d_G(u)$. Thus there are three types of reverse edges of follows:

$$\begin{aligned} CE_{44} &= \{uv \in E(G) | c_u = c_v = 4\}, & |CE_{44}| &= 6n. \\ CE_{41} &= \{uv \in E(G) | c_u = 4, c_v = 1\}, & |CE_{41}| &= 18n^2 + 6n. \\ CE_{11} &= \{uv \in E(G) | c_u = c_v = 1\}, & |CE_{11}| &= 18n^2 - 12n. \end{aligned}$$

In the following theorem, we compute the geometric-arithmetic reverse index of silicate networks.

Theorem 2.1. *The geometric-arithmetic reverse index of silicate networks is given by*

$$GAC(SL_n) = \frac{162}{5}n^2 - \frac{6}{5}n.$$

Proof. By definition, we have

$$GAC(G) = \sum_{uv \in E(G)} \frac{2\sqrt{c_u c_v}}{c_u + c_v}$$

Thus

$$\begin{aligned} GAC(SL_n) &= \sum_{CE_{44}} \frac{2\sqrt{c_u c_v}}{c_u + c_v} + \sum_{CE_{41}} \frac{2\sqrt{c_u c_v}}{c_u + c_v} + \sum_{CE_{11}} \frac{2\sqrt{c_u c_v}}{c_u + c_v} \\ &= \left(\frac{2\sqrt{4 \times 4}}{4 + 4} \right) 6n + \left(\frac{2\sqrt{4 \times 1}}{4 + 1} \right) (18n^2 + 6n) + \left(\frac{2\sqrt{1 \times 1}}{1 + 1} \right) (18n^2 - 12n) \\ &= \frac{162}{5}n^2 - \frac{6}{5}n. \end{aligned}$$

□

In the following theorem, we compute the sum connectivity reverse index of silicate networks.

Theorem 2.2. *The sum connectivity reverse index of silicate networks is given by*

$$SC(SL_n) = \left(\frac{18}{\sqrt{5}} + \frac{18}{\sqrt{2}} \right) n^2 + \left(\frac{6}{\sqrt{5}} - \frac{9}{\sqrt{2}} \right) n.$$

Proof. By definition, we have

$$SC(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{c_u + c_v}}.$$

Thus

$$\begin{aligned} SC(SL_n) &= \sum_{CE_{44}} \frac{1}{\sqrt{c_u + c_v}} + \sum_{CE_{41}} \frac{1}{\sqrt{c_u + c_v}} + \sum_{CE_{11}} \frac{1}{\sqrt{c_u + c_v}} \\ &= \left(\frac{1}{\sqrt{4+4}} \right) 6n + \left(\frac{1}{\sqrt{4+1}} \right) (18n^2 + 6n) + \left(\frac{1}{\sqrt{1+1}} \right) (18n^2 - 12n) \\ &= \left(\frac{18}{\sqrt{5}} + \frac{18}{\sqrt{2}} \right) n^2 + \left(\frac{6}{\sqrt{5}} - \frac{9}{\sqrt{2}} \right) n. \end{aligned}$$

□

3. Results for Hexagonal Networks

A triangular tiling is used in the construction of hexagonal networks. A hexagonal network is symbolized by HX_n where n is the number of vertices in each side of hexagon. A hexagonal network of dimension six is depicted in Figure 2.




Figure 2: A 6-dimensional hexagonal network

Let H be the graph of hexagonal network HX_n . By calculation, we obtain that $|V(HX_n)| = 3n^2 - 3n + 1$ and $|E(HX_n)| = 9n^2 - 15n + 6$. From Figure 2, one can see that the vertices of HX_n are either of degree 3, 4 or 6. Then $\Delta(H) = 6$. In H , by algebraic method, there are five types of edges based on the degree of the end vertices of each edge as follows:

$$E_{34} = \{uv \in E(H) | d_H(u) = 3, d_H(v) = 4\}, \quad |E_{34}| = 12.$$

$$E_{36} = \{uv \in E(H) | d_H(u) = 3, d_H(v) = 6\}, \quad |E_{36}| = 6.$$

$$E_{44} = \{uv \in E(H) | d_H(u) = d_H(v) = 4\}, \quad |E_{44}| = 6n - 18.$$

$$E_{46} = \{uv \in E(H) | d_G(u) = 4, d_G(v) = 6\}, \quad |E_{46}| = 12n - 24.$$

$$E_{66} = \{uv \in E(H) | d_G(u) = d_G(v) = 6\}, \quad |E_{66}| = 9n^2 - 33n + 30.$$

Clearly, we have $c_u = \Delta(H) - d_H(u) + 1 = 7 - d_H(u)$. Thus there are five types of reverse edges as follows:

$$\begin{aligned} CE_{43} &= \{uv \in E(H) | c_u = 4, c_v = 3\}, & |CE_{43}| &= 12. \\ CE_{41} &= \{uv \in E(H) | c_u = 4, c_v = 1\}, & |CE_{41}| &= 6. \\ CE_{33} &= \{uv \in E(H) | c_u = c_v = 3\}, & |CE_{33}| &= 6n - 18. \\ CE_{31} &= \{uv \in E(H) | c_u = 3, c_v = 1\}, & |CE_{31}| &= 12n - 24. \\ CE_{11} &= \{uv \in E(H) | c_u = c_v = 1\}, & |CE_{11}| &= 9n^2 - 33n + 30. \end{aligned}$$

In the following theorem, we compute the geometric-arithmetic reverse index of hexagonal networks.

Theorem 3.1. *The geometric-arithmetic reverse index of hexagonal networks is given by*

$$GAC(HX_n) = 9n^2 + (6\sqrt{3} - 27)n + \left(\frac{48\sqrt{3}}{7} + \frac{24}{5} + 12\right).$$

Proof. By definition, we have

$$GAC(H) = \sum_{uv \in E(H)} \frac{2\sqrt{c_u c_v}}{c_u + c_v}.$$

Thus

$$\begin{aligned} GAC(HX_n) &= \sum_{CE_{43}} \frac{2\sqrt{c_u c_v}}{c_u + c_v} + \sum_{CE_{41}} \frac{2\sqrt{c_u c_v}}{c_u + c_v} + \sum_{CE_{33}} \frac{2\sqrt{c_u c_v}}{c_u + c_v} + \sum_{CE_{31}} \frac{2\sqrt{c_u c_v}}{c_u + c_v} + \sum_{CE_{11}} \frac{2\sqrt{c_u c_v}}{c_u + c_v} \\ &= \left(\frac{2\sqrt{4 \times 3}}{4 + 3}\right) 12 + \left(\frac{2\sqrt{4 \times 1}}{4 + 1}\right) 6 + \left(\frac{2\sqrt{3 \times 3}}{3 + 3}\right) (6n - 18) \\ &\quad + \left(\frac{2\sqrt{3 \times 1}}{3 + 1}\right) (12n - 24) + \left(\frac{2\sqrt{1 \times 1}}{1 + 1}\right) (9n^2 - 33n + 30) \\ &= 9n^2 + (6\sqrt{3} - 27)n + \left(\frac{48\sqrt{3}}{7} + \frac{24}{5} + 12\right). \end{aligned}$$

□

In the following theorem, we compute the sum connectivity reverse index of hexagonal networks.

Theorem 3.2. *The sum connectivity reverse index of hexagonal networks is given by*

$$SC(HX_n) = \frac{9}{\sqrt{2}}n^2 + \left(\sqrt{6} + 6 - \frac{33}{\sqrt{2}}\right)n + \left(\frac{12}{\sqrt{7}} + \frac{6}{\sqrt{5}} - 3\sqrt{6} - 12 + \frac{30}{\sqrt{2}}\right).$$

Proof. By definition, we have

$$SC(H) = \sum_{uv \in E(H)} \frac{1}{\sqrt{c_u + c_v}}.$$

Thus

$$\begin{aligned} SC(HX_n) &= \sum_{CE_{43}} \frac{1}{\sqrt{c_u + c_v}} + \sum_{CE_{41}} \frac{1}{\sqrt{c_u + c_v}} + \sum_{CE_{33}} \frac{1}{\sqrt{c_u + c_v}} + \sum_{CE_{31}} \frac{1}{\sqrt{c_u + c_v}} + \sum_{CE_{11}} \frac{1}{\sqrt{c_u + c_v}} \\ &= \left(\frac{1}{\sqrt{4 + 3}}\right) 12 + \left(\frac{1}{\sqrt{4 + 1}}\right) 6 + \left(\frac{1}{\sqrt{3 + 3}}\right) (6n - 18) \\ &\quad + \left(\frac{1}{\sqrt{3 + 1}}\right) (12n - 24) + \left(\frac{1}{\sqrt{1 + 1}}\right) (9n^2 - 33n + 30) \\ &= \frac{9}{\sqrt{2}}n^2 + \left(\sqrt{6} + 6 - \frac{33}{\sqrt{3}}\right)n + \left(\frac{12}{\sqrt{7}} + \frac{6}{\sqrt{5}} - 3\sqrt{6} - 12 + \frac{30}{\sqrt{2}}\right). \end{aligned}$$

□

References

-
- [1] V.R.Kulli, *College Graph Theory*, Vishwa International Publications, Gulbarga, India, (2012).
- [2] V.R.Kulli, *Atom bond connectivity reverse and product connectivity reverse indices of oxide and honeycomb networks*, submitted.
- [3] V.R.Kulli, *On the sum connectivity reverse index of oxide and honeycomb networks*, Journal of Computer and Mathematical Sciences, 8(9)(2017), 408-413.
- [4] S.Ediz, *Maximal graphs of the first reverse Zagreb beta index*, TWMS J. App. Eng. Math. accepted for publication.
- [5] V.R.Kulli, *On the product connectivity reverse index of silicate and hexagonal networks*, International Journal of Mathematics and its Applications, 5(4B)(2017), 175179.
- [6] V.R.Kulli, *Reverse Zagreb and reverse hyper-Zagreb indices and their polynomials of rhombus silicate networks*, submitted.
- [7] I.Gutman and N.Trinajstic, *Graph theory and molecular orbitals, Total π -electron energy of alternant hydrocarbons*, Chem. Phys. Lett., 17(1972), 535-538.
- [8] V.R.Kulli, *K-Banhatti indices of graphs*, Journal of Computer and Mathematical Sciences, 7(4)(2016), 213-218.
- [9] V.R.Kulli, *On K-hyper-Banhatti indices and coindices of graphs*, International Research Journal of Pure Algebra, 6(5)(2016), 300-304.
- [10] V.R.Kulli, *New K Banhatti topological indices*, International Journal of Fuzzy Mathematical Archive, 12(1)(2017), 29-37.
- [11] V.R.Kulli, *The Gourava indices and coindices of graphs*, Annals of Pure and Applied Mathematics, 14(1)(2017), 33-38.
- [12] V.R.Kulli, *Computation of general topological indices for titania nanotubes*, International Journal of Mathematical Archive, 7(12)(2016), 33-38.
- [13] V.R.Kulli, *First multiplicative K Banhatti index and coindex of graphs*, Annals of Pure and Applied Mathematics, 11(2)(2016), 79-82.
- [14] V.R.Kulli, *Some new multiplicative geometric-arithmetic indices*, Journal of Ultra Scientist of Physical Sciences A, 29(2)(2017), 52-57.
- [15] V.R.Kulli, *General topological indices of tetrameric 1,3-adamantane*, International Journal of Current Research in Science and Technology, 3(8)(2017), 26-33.
- [16] V.R.Kulli, *Computing topological indices of dendrimer nanostars*, International Journal of Mathematics and its Applications, 5(3-B)(2017), 163-169.
- [17] I.Gutman, V.R.Kulli, B.Chaluvaraju and H.S.Baregowda, *On Banhatti and Zagreb indices*, Journal of the International Mathematical Virtual Institute, 7(2017), 53-67.
- [18] V.R.Kulli, *Computation of some topological indices of certain networks*, International Journal of Mathematical Archive, 8(2)(2017), 99-106.